Original Research Article

DOI: http://dx.doi.org/10.18203/2394-6040.ijcmph20172179

Prevalence and risk factors of urinary incontinence during antenatal period in women delivering in a tertiary care center of Northern India

Bhanu Priya¹*, Nilanchali Singh², Shalini Rajaram¹

Department of Obstetrics and Gynecology, Guru Teg Bahadur Hospital, New Delhi, Maulana Azad Medical Hospital, Delhi, India

Received: 10 April 2017 **Revised:** 01 May 2017 **Accepted:** 04 May 2017

*Correspondence: Dr. Bhanu Priya,

E-mail: drpriyabhanu@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Urinary incontinence (UI) is a commonly encountered entity in antenatal and postpartum period. It is known to have detrimental effects on quality of life in approximately 54.3% of all pregnant women. However, the true prevalence of UI is still not known, especially in the South East Asia. This study has investigated the prevalence and risk factors of urinary incontinence during antenatal period.

Methods: 500 women admitted in postnatal ward were interviewed retrospectively about the UI occurred during antenatal period, using questionnaire. The study was done over a period of six months. Detailed information regarding risk factors of urinary incontinence was included in questionnaire based on previous validated studies. Data was analyzed using SPSS 20 using chi square test. P value <0.05 was taken as significant.

Results: Total 500 women were interviewed and based on questionnaire. 202 reported incontinence, thus giving a prevalence of 40.4%, out of that 192 (45.5%) had stress urinary incontinence (SUI), 40(19.8%) had urge urinary incontinence (UUI) and 70 (34.6%) had mixed incontinence. The risk factors like bladder infections and chronic cough had significant association with urinary incontinence. Advancing gestation had worsening effects on the symptoms of UI (p =0.000).

Conclusions: Urinary incontinence is a common and neglected problem with poor treatment seeking behavior. SUI is more common in pregnant women than urge or mixed incontinence. Obstetricians need to ask have precise interrogation regarding UI in pregnant women to diagnose and treat the problem.

Keywords: Prevalence, Risk factors, Urinary incontinence, Questionnaire, Treatment seeking behavior

INTRODUCTION

Urinary incontinence (UI) is a significant health concern and has been shown to impair women's activities and also affects the physical and mental health. UI is higher in few specific groups like pregnancy and have been reported a prevalence of 32%-64% for all UI and 59% for stress urinary incontinence (SUI) and 40% for mixed incontinence. Women with UI have poorer treatment seeking behavior for UI. Approximately 54.3% of all pregnant women have detrimental effects on quality of life. Pregnancy is an independent risk factor for UI that

leads to weakened pelvic floor muscle because of pregnancy related physiological and anatomical changes. However, despite the multiple references in literature on effects of, parity, age of women, pathophysiological changes in pregnancy and mode of delivery, on the pelvic floor and lower urinary tract symptoms remain inconclusive.⁴

Prenatal physiological changes such as increasing pressure of the advancing gravid uterus and growing fetal weight on pelvic floor muscle (PFM) throughout the pregnancy, along with the pregnancy-related changing

levels of hormones such as progesterone, estrogen, and relaxin, may lead to reduced strength and sphincteric function of PFM.⁵⁻⁸ PFM weakness causes bladder-neck and urethral hypermobility, leading to urethral sphincter incompetence. Hence, when intra-abdominal pressure is increased with coughing, sneezing, laughing, or moving, the pressure inside the bladder exceeds the urethral closure pressure and the urethral sphincter is not strong enough to maintain urethral closure. Urinary leakage will be the result. SUI is common during pregnancy and puerperium. After delivery, SUI symptoms resolve in the vast majority of cases.⁹ The healing process may take some time after the delivery, but in a significant percentage of women, it can persist in subsequent stages of life. 10 In primipara women, SUI symptoms tend to resolve within 3 month after delivery.¹¹

Multiple risk factors have been associated with UI. Significant risk factors for UI in pregnancy were maternal age ≥35 years, body mass index and parity. ¹² Smoking, diabetes, chronic cough and high intake of caffeine are identifiable risk factors for UI. ¹³⁻¹⁵

The information available on UI in postpartum women from India is sparse. Therefore, a study on UI is designed which would add information regarding prevalence of UI in this part of world as incidence and prevalence estimates of incontinence vary widely. Only a few population based studies have investigated prevalence of urinary incontinence during pregnancy by type and severity. Also, data on risk factors for incontinence in pregnancy are scarce.

METHODS

This observational study was carried out over six weeks in the department of obstetrics and gynecology, at a tertiary care hospital in Northern India. All the healthy postpartum women admitted in the post natal ward, who had singleton pregnancy, were included in the study. Total 500 subjects were included in the study. Informed consent was obtained from the participants prior to conducting the study and the women were interviewed about the given questionnaire. The study was done over a period of six months from March 2016 to September 2016. Urinary incontinence was assessed using standardized questions based on questions validated in previous studies. 17,18 The questionnaire covered the risk factors, symptoms and severity of the UI. Women who had leakage of urine at least once a month, regardless of amount were defined as UI. Women were defined as having stress urinary incontinence if leakage of urine was associated with coughing, laughing, sneezing or physical activity, urge incontinence, if leakage was preceded or accompanied by urgency, and mixed incontinence when symptoms of both stress (SUI) and urge (UUI). These definitions follow the standardized terminology for lower urinary tract symptoms endorsed by the International Continence Society. 19 Severity of UI was assessed as slight, moderate or severe based on the frequency and amount of leakage using the incontinence severity index. 17,18 The questionnaire was used to gather information about socio-demographic pattern, predisposing and risk factors, treatment seeking behavior including the prevalence of UI.

Data analysis was done using SPSS version 20. Odd's ratio and chi- square test was used for univariate analysis.

RESULTS

During the study, 500 women were interviewed. The mean age of study population was 25.1 years with a range of 19-40 years. The demographic characteristics like age, weight, height, BMI, parity and past delivery mode were not found to have a significant association with UI (Table 1).

Table 1: Study group characteristics.

Characteristics	Group I (n=202) (incontinence present)	Group II (n=298) (incontinence absent)	P value
Age (years)	25.1±3.2	24.9±3.6	0.703
Weight (kg)	57.7±5.0	57.6±5.1	0.912
Height (cm)	151.7±5.0	15 1.9±5.2	0.815
BMI (kg/m ²)	24.9±2.3	25.0±2.1	0.886
Parity			
Primipara	81	104	0.294
Multipara	163	152	
Past delivery mode			
No past delivery	91	86	0.535
Previous normal delivery	126	88	
Previous cesarean section	76	33	
Bladder habits alteration	371	129	0.187
Caffeine intake	203	297	1.000
Bladder infection	40	460	0.013
Pelvic pain	35	465	0.405
Chronic cough	48	452	0.001
Constipation	11	489	0.15

Table 2: Risk factors for urinary incontinence.

Risk factors	Odd's ratio	95% confidence interval
Bladder infection	3.75	1.2-11.2
Pelvic pain	1.58	0.5-4.7
Chronic cough	4.99	1.7-14.5
Constipation	3.23	0.7-13.3




Figure 1: Bar diagram of frequency of urinary incontinence and treatment seeking behavior.

Out of 500 participants, 185 primipara participated in study and of those 185 participants, 81 subjects reported incontinence. The prevalence of incontinence was 43.7% in primipara. Similarly, 315 multipara took part in the study and 163 had incontinence which gives a prevalence of 53.4% in multipara women.

In the study population chronic cough (p=0.001) and bladder infections (p=0.013) had significant association with UI. Table 2 shows p value and odds ratio for the risk factors.

Out of 500 women 202 reported incontinence, thus giving a prevalence of 40.4%, out of that 192 (45.5%) had SUI, 40(19.8%) had UUI and 70 (34.6%) had mixed incontinence. Among the incontinent women 31/202 (15.3%) sought treatment (Figure 1). The treatment seeking behavior was not related to severity of UI.

DISCUSSION

Pregnancy is associated with lower urinary tract symptoms and UI is also very troublesome complaint. The most common type of UI is SUI. Subjects who had incontinence in our study was 40.4% which is in the range of the previous study in the literateure. In our study SUI was the most common form of incontinence (45.5%) followed by mixed incontinence (34.6%) and thereafter UUI (19.8%). This was also in accordance to previous studies in the literature, which shows that SUI is the most common form of UI suggests that prenatal pelvic floor exercise can be effective in prevention of SUI during pregnancy and postpartum period. It has been observed in the previous studies that six weeks pelvic

floor muscle exercise programme was able to decrease the severity of symptoms of incontinence with SUI.

Despite that nearly half of the pregnant women population was affected by UI, the treatment seeking behavior was only 15.3% of all incontinent women. It was also very interesting to note that only 6.9% of women with very severe UI did seek treatment.

The prevalence of incontinence in primipara was slightly higher than previously reported in literature.²¹ In our study 53.4% multipara had incontinence which shows that parity has association with incontinence but it was not stastically significant.

In this study, sociodemographic pattern did not have any association with UI. Chronic cough and bladder infections had significant relation with UI. After univariate analysis the Odd's ratio for UI in subjects with chronic cough and bladder infections was 4-5 times more. Chronic cough itself has direct association with UI. 24 Sudden increase in intra-abdominal pressure leads to exhaustion of pelvic floor muscles. Hence, UI was found to be more common with multiparity, chronic cough and bladder infections in our study.

The study population was limited to our tertiary care center. Also, none of the subjects was examined gynaecologically for UI and UI was only inquired verbally which may reduce the accuracy of diagnosis.

CONCLUSION

A significant number of antenatal women are suffering from UI and it is substantially affecting the quality of life. The women in India are suffering from unawareness about the condition, hence making it difficult to diagnose for physician and further early treatment. Primary prevention of UI is needed by increasing public awareness about pelvic floor muscle exercise during and after pregnancy.

Funding: No funding sources Conflict of interest: None declared

Ethical approval: The study was approved by the

Institutional Ethics Committee

REFERENCES

1. Handa VL, Zyczynski HM, Burgio KL, Fitzgerald MP, Borello-France D, Janz NK, et al. The impact

- of fecal and urinary incontinence on quality of life 6 months after childbirth. Am J Obstet Gynecol. 2007;636:1–6.
- 2. Milsom I, Altman D, Lapitan MC, Nelson R, Sillén U, Thom D. Epidemiology of urinary (UI) and faecal (FI) incontinence and pelvic organ prolapse (POP). In: Abrams P, Cardozo L, Khoury S, Wein A (editors). Incontinence, 4th ed. Committee 1. Paris, France: Health Publication Ltd; 2009: 37-111.
- Dolan LM, Walsh D, Hamilton S, Marshall K, Thompson K, Ashe RG. A study of quality of life in primigravidae with urinary incontinence. Int Urogynecol J Pelvic Floor Dysfunct. 2004;15:160– 4.
- 4. Koebl H, Nitti V, Baessler K, Salvatore S, Sultan A, Yamaguchi O. Pathophysiology of urinary incontinence, fecal incontinence and pelvic organ prolapse. In: Abrams P, Cardozo L, Khoury, Wein A (editors), Incontinence. 4th ed. Plymouth: Health Publication Ltd; 2009: 255-330.
- 5. Davis K, Kumar D. Pelvic floor dysfunction: a conceptual framework for collaborative patient-centred care. J Adv Nurs. 2003;43(6):555–68.
- Morkved S, Bo K, Schei B, Salvesen KA. Pelvic floor muscle training during pregnancy to prevent urinary incontinence: a singleblind randomized controlled trial. Obstet Gynecol. 2003;101(2):313– 9.
- 7. Hilton P, Dolan LM. Pathophysiology of urinary incontinence and pelvic organ prolapse. Br J Obstet Gynaecol. 2004;111(1):5–9.
- 8. Kristiansson P, Samuelsson E, Schoultz B, Svardsudd K. Reproductive hormones and stress urinary incontinence in pregnancy. Acta Obstet Gynecol Scand. 2001;80:1125–30.
- 9. Viktrup L, Rortveit G, Lose G. Risk of stress urinary incontinence twelve years after the first pregnancy and delivery. Obstet Gynecol. 2006;108(2):248–54.
- 10. Thom DH, Rortveit G. Prevalence of postpartum urinary incontinence: a systematic review. Acta Obstet Gynecol Scand. 2010;89(12):1511–22.
- 11. Viktrup L, Lose G, Rolf M, Barfoed K. The frequency of urinary symptoms during pregnancy and puerperium in the primipara. Int Urogynecol J. 1993;4(1):27–30.
- 12. Cerruto MA, D'Elia C, Aloisi A, Fabrello M, Artibani W. Prevalence, Incidence and Obstetric Factors' Impact on Female Urinary Incontinence in Europe: A Systematic Review. Urol Int. 2013;90(1):1-9.
- 13. Kirss F, Lang K, Toompere K, Veerus P. Prevalence and risk factors of urinary incontinence among Estonian postmenopausal women. Springer Plus. 2013,2:524.
- Swanson JG, Kaczorowski J, Skelly J, Finkelstein M. Urinary incontinence: common problem among

- women over 45. Can Fam Physician. 2005,51(1):84–5.
- Gleason JL, Richter HE, Redden DT, Goode PS, Burgio KL, Markland AD. Caffeine and urinary incontinence in US women. Int Urogynecol J. 2013;24(2):295–302.
- Burgio KL, Locher JL, Zyczynski H, Hardin JM, Singh K. Urinary incontinence during pregnancy in a racially mixed sample: characteristics and predisposing factors. Int Urogynecol J Pelvic Floor Dysfunct. 1996;7:69-73.
- 17. Sandvik H, Hunskaar S, Seim A, Hermstad R, Vanvik A and Bratt H. Validation of a severity index in female urinary incontinence and its implementation in an epidemiological survey. J Epidemiol Community Health. 1993;47:497-9.
- Sandvik H, Seim A, Vanvik A and Hunskaar S. A severity index for epidemiological surveys of female urninary incontinence: Comparison with 48-hour Pad-Weighing Tests. Neurourol Urodynamics. 2000;19:137-45.
- Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, et al. The standardisation of terminology in lower urinary tract function: report from the standardisation sub-committee of the International Continence Society. Urol. 2003;61:37– 49.
- 20. Bø K, Pauck Øglund G, Sletner L, Mørkrid K, Jenum AK. The prevalence of urinary incontinence in pregnancy among a multi-ethnic population resident in Norway. BJOG. 2012;119(11):1354-60.
- 21. Zhu L, Li L, Lang JH, Xu T. Prevalence and risk factors for peri and postpartum urinary incontinence in primiparous women in China: a prospective longitudinal study. Int Urogynecol J. 2012;23(5):563-72.
- 22. Hem L, Qamar-ur-Nisa, Jabeen A, Pushpa G; The Pattern of Urinary Tract Symptoms During Pregnancy. Pak J Med Sci. 2011;27(4):827-31.
- 23. Sangsawang B, Serisathien Y. Effect of pelvic floor muscle exercise programme on stress urinary incontinence among pregnant women. J Adv Nurs. 2012;68(9):1997-2007.
- 24. Perera J, Kirthinanda DS, Wijeratne S, Wickramarachchi TK. Descriptive cross sectional study on prevalence, perceptions, predisposing factors and health seeking behaviour of women with stress urinary incontinence. BMC Women's Health. 2014;14:78.

Cite this article as: Priya B, Singh N, Rajaram S. Prevalence and risk factors of urinary incontinence during antenatal period in women delivering in a tertiary care center of Northern India. Int J Community Med Public Health 2017;4:2071-4.