

## Original Research Article

# Factors associated with level of birth preparedness among pregnant mothers at Kerugoya county referral hospital

Olive W. Karimi<sup>1</sup>, Mary W. Murigi<sup>2\*</sup>, Anne Pertet<sup>3</sup>, Careena O. Odawa<sup>3</sup>

<sup>1</sup>Department of Nursing, Kerugoya County Referral Hospital, Kirinyaga County, Kenya

<sup>2</sup>School of Health Science, Kirinyaga University, Kirinyaga County, Kenya

<sup>3</sup>Department of Community Health and Development, Great Lakes University of Kisumu, Kisumu County, Kenya

**Received:** 24 August 2017

**Accepted:** 23 September 2017

### \*Correspondence:

Dr. Mary W. Murigi,

E-mail: [mmurigi@kyu.ac.ke](mailto:mmurigi@kyu.ac.ke)

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background:** Birth preparedness is the advance preparation made by an expectant mother which ensures access to skilled care. In Africa, the risk of pregnancy related deaths is 300 times more than in the developed world. In Kenya, out of 10 expectant mothers who access antenatal care services only 4 deliver in a health facility.

**Methods:** This was a descriptive cross sectional study conducted at outpatient Maternal Child Health and Family Planning Clinic at Kerugoya County Hospital. The study utilized convenience sampling technique to determine the study population. The research instruments were an In-depth interview guide and semi-structured questionnaires. Data was managed using SPSS and analysis done using descriptive statistics and Chi-square tests. Statistical significance was set at  $p < 0.05$ .

**Results:** A sample of 202 women participated in the study. 47.5% of the participants were adequately prepared for birth. Having a higher level of education was significantly associated with birth preparedness ( $p = 0.021$ ). The number of children per woman had a significant influence on level of birth preparedness with women who had no children less likely to be prepared for birth compared to those with one or more children ( $p = 0.002$ ). Clients who attended Antenatal Care (ANC) at least 3 times were prepared for birth compared to those who visited either once or twice ( $p = 0.027$ ).

**Conclusions:** Overall, women of reproductive age lack birth preparedness. There is therefore need to enhance birth preparedness awareness campaigns at ANC visits targeting women in their third trimester.

**Keywords:** Birth preparedness, Pregnant women, Third trimester

## INTRODUCTION

Birth Preparedness is the process of preparing for a normal birth during pregnancy or emergency and the actions required to be taken in case of complications.<sup>1</sup> According to the Integrated Management of Pregnancy and Childbirth (IMPAC), it is a plan made by expectant mothers and should be discussed and reviewed by a skilled attendant during each Ante Natal visit and

especially one month prior to the expected date of delivery.<sup>2</sup>

Focused antenatal care (FANC) was endorsed by WHO as one of the pillars of safe motherhood initiative (SMI) which is a global strategy to be used as a policy guideline in all facilities that offers Maternal Child Health (MCH).<sup>2</sup> The strategy was to reduce three major delays during childbirth that contribute to both maternal and child mortality and morbidity especially in developing countries.<sup>3</sup> The three delays are: delay in deciding to seek

medical care, delay in reaching the health facility and delay in receiving care at the health facility. Advance birth preparedness is therefore required to minimize these delays and ensure all mothers receive skilled health care during and after childbirth.<sup>4</sup>

In Kenya, according to the Kenya Demographic Health Survey (KDHS)2008/09, the maternal mortality rate was at 488/100,000 live births with 42% of mothers accessing skilled deliveries which means that over 50% of mothers and newborns are in danger should any complication occur during delivery or after.<sup>5</sup> By the year 2015 maternal deaths according to the millennium development goals were to be reduced to about 147 /100,000 live births. According to KDHS 2014/15, Maternal mortality is still unacceptably high at 362/1000 live births in Kenya.<sup>6</sup> According to the National Road Map of 2010 in the Ministry of Health in Kenya over 50% of deliveries in the country among Kenyan women are attended by unskilled persons.<sup>7</sup> Also 92% of pregnant women access ANC services but only 43% deliver in a health facility which is below the target of 90% by 2015.<sup>7</sup>

The reasons for the non-improvement in maternal health are delays in decision making and seeking care when problems occur. Birth preparedness is aimed to assist expectant mothers to have enough resources for use during the birth process and to access timely skilled care at birth.<sup>8</sup> Lack of advance planning for childbirth is one of the contributing factors why pregnant mothers delay in seeking and reaching the health care facilities. There is considerable time for this preparation as the duration of pregnancy is long.

Pregnancy and childbirth complications in Kerugoya County Referral Hospital, Kenya have been a challenge. Despite the high ANC attendance of 90%, mothers still arrive in maternity in second stage of labour resulting to maternal complications. No study has been done in Kirinyaga County on the level of birth preparedness. Thus the findings from this study will contribute to the existing knowledge on birth preparedness and generate further discussions on how to make pregnancy safe at the same time improve on the maternal and newborn health in the County and the country at large.

## METHODS

The study was a descriptive cross sectional survey conducted at outpatient Maternal Child Health and Family Planning Clinic (MCH/FP) at Kerugoya County referral Hospital. The study took place between the months of August 2015 to April 2016. The study population was pregnant mothers attending ANC at Kerugoya County Referral. The unit of analysis was those pregnant mothers who had reached 28 weeks and had attended the clinic at least once previously. The study utilized convenience sampling. The sampling strategy was to sample to saturation. Any eligible woman who visited the health facility and consented was enrolled until the researcher achieved a sample size of 212

women. The research instruments were an In-depth interview guide and self-administered semi-structured questionnaires. Quantitative data analysis was conducted using statistical analysis for quantitative data was performed using IBM Statistical Package for Social Sciences (SPSS). The data analysis involved descriptive statistics and Chi-square tests. Descriptive statistics were used to describe the demographic characteristics, while Chi-square test was used to show the relationship between demographic and socioeconomic characteristics and level of birth preparedness. Statistical significance was set at  $p < 0.05$ .

## RESULTS

### *Socio demographic characteristics of the participants*

A total of 212 women took part in the study. The mean age of the study subjects was 26.1 ( $\pm 5.6$  SD) years. The majority 61.9% of the study participants were of the age group 20-30 years. Most of the respondents 57.1% only had a primary level education which is the basic education in Kenya. 75.9% of the women were married and majority had between one to two children 66.9% (Table 1).

**Table 1: Socio- demographic characteristics.**

Variable	Frequency	Percentage (%)
Age	less than 20 years	12.4
	20 to 30 years	61.9
	over 30 years	25.7
Religion	Protestant	64.4
	Catholic	33.7
	Muslim	2.0
Education	Tertiary	5.8
	Secondary	26.7
	Primary	67.5
Occupation	Housewife	4.5
	Farming	62.9
	Business	6.4
	Salaried	21.8
	Other	4.5
Marital status	Single	4.5
	Married	75.7
	Other	2.0
Number of children	0	30.7
	1	34.2
	2	32.7
	3	2.5
Number of ANC visits	<4 Visits	52.0
	$\geq 4$ Visits	48.0

### *Level of birth preparedness*

An overall composite level of preparedness was classified into two, prepared or not prepared. If any combination of at least three positive actions were mentioned, this was

referred to as ‘prepared’, otherwise preparedness was classified as ‘not prepared’. Results showed that overall 47.5% were prepared for birth while majority were not prepared. All women were prepared by saving some money and all of them had identified place of delivery (Figure 1).




Figure 1: Level of birth preparedness.

**Association between demographic factors and birth preparedness**

A woman’s age was significantly associated with birth preparedness. Younger women aged less than 25 years

were 1.4 times likely to be birth prepared compared to those aged above 25 years. Noteworthy is that Marital status of the pregnant women had no association with birth preparedness (p=0.736) (Table 2).

**Association between socio economic factors and birth preparedness**

Majority of the respondents 57.43% had primary education while 26.7% had secondary school education. Of those who had secondary education, 96.3% had birth preparedness compared to those who had primary education. Religion had no influence on birth preparedness. However the number of children the women had, had a significant influence on level of birth preparedness. Women who had no children were less likely to be prepared for birth compared to those with one child (OR=0.4; 95%CI= [0.2-0.8] or 2 children (OR=0.3; 95%CI= [0.1-0.6]). The number of ANC visits was associated with birth preparedness. Of those who attended ANC at least 3 times or more, 51.6% of them were prepared for birth compared to 32.6% of those prepared but only visited ANC either once or twice (p=0.027).

Table 2: Association between demographic factors and birth preparedness.

	Overall (N=202)	Birth preparedness		Odds ratio (95% CI)	P value
	N (%)	Yes N (%)	No N (%)		
<b>Age</b>					
25 years and below	97 (48.01)	55 (56.7)	42 (43.3)	1.4[1.1-3.6]	0.012
>25 years	105 (51.98)	41 (39.05)	64 (60.95)		
<b>Marital status</b>					
Single	45 (22.73)	20 (44.44)	25 (55.56)	0.9[0.6-1.3]	0.736
Married	153 (77.27)	73 (47.71)	80 (52.29)		




Figure 2: Perception of women on birth preparedness.

**Table 1: Association between socio economic factors and birth preparedness.**

	Overall (N=202)	Birth preparedness		Odds ratio (95% CI)	P value
	Overall	Yes	No		
	N (%)	N (%)	N (%)		
<b>Education level</b>					
Tertiary	32 (15.84)	32 (100)	0 (0)	NA	0.993
Secondary	54 (26.73)	52 (96.30)	2 (3.7)		0.021
Primary	116 (57.43)	12 (10.34)	104 (89.66)	Ref	-
<b>Religion</b>					
Protestant	130 (64.36)	60 (46.15)	70 (53.85)	Ref	-
Catholic	68 (33.36)	34 (50.0)	34 (50)	0.9[0.5-1.5]	0.607
Muslim	4 (1.98)	2 (50.0)	2 (50.0)	0.9[0.1-6.3]	0.879
<b>Number of children</b>					
0	62 (30.69)	19 (30.65)	43 (69.35)	Ref	-
1	69 (34.16)	36 (52.17)	33 (47.83)	0.4[0.2-0.8]	0.014
2	66 (32.67)	39 (59.09)	27 (40.91)	0.3[0.1-0.6]	0.002
3	5 (2.48)	2 (40.0)	3 (60.0)	0.7[0.1-4.3]	0.667
<b>ANC visits</b>					
1-2	43 (21.29)	14 (32.56)	29 (67.44)	0.6[0.4-0.9]	0.027
3 and above	159 (78.71)	82 (51.57)	77 (48.43)		

**Perception of women on birth preparedness**

The perception was assessed using a Likert-scale questionnaire. Six indicators of women perception of birth preparedness were used in this analysis. The likert scale values were as follows; strongly disagree (-2), disagree (-1), neutral/don't know (0), agree (1) and strongly agree was coded (2). Mean of the 6 indicators was calculated as scores. Overall, amongst the 202 women mean perception score was found to be (-2.93) ±2.35 SD). The distribution given in Figure 2 indicated that in all the six indicators the perception towards birth preparedness was negative and those with positive perception were below 5%.

**DISCUSSION**

Overall, majority 52.5% of the women were not prepared for birth. During the in depth interview the women were found to save money through women groups which apparently was the norm in the rural set up. In addition to this although all the mothers (100%) attempted to save money mostly to 'buy clothes for baby' and 'cotton wool', transport arrangements did not seem to appear as part of birth preparedness and this contributes to the delay in reaching the health facility as time is wasted as means of transport is sought. This concurs with the report by the National Road Map on Maternal and Newborn Health 2010 that one of the reasons for non-improvement of maternal and newborn indicators in Kenya is delay in reaching the health facility.<sup>7</sup>

This study established that the age of the women was significantly associated with birth preparedness. Younger

women aged 25 years and below were 1.4 times more likely to be birth prepared compared to those aged 25 years and above. This is consistent with a study on factors associated with the use of maternal health services in southern Nigeria which showed younger mothers were more likely to prepare for childbirth than their elder counterparts.<sup>9</sup> Yet another study done in southern Ethiopia titled birth preparedness and complication readiness also indicated that mothers who were aged 25 years and below were more likely to prepare for birth than their older counterparts.<sup>10</sup>

The study established that the number of children the women had was significantly associated with level of birth preparedness. Women who had no children were less likely to be prepared for birth compared to those with one child or two children. A study conducted in urban slums in Kenya found that the likelihood of preparing and delivering at a health facility significantly decreased as parity increases.

Results from this study also established that women who had secondary education, 96.3% had birth preparedness compared to 10.34% of those who had primary education. From the in-depth interviews, education enabled them to understand the importance of birth preparedness as they could also get information from other sources as magazines since they can read. A woman who is educated is able to make wise decisions about her own health than one who is not.<sup>11</sup> The status of women is likely to be enhanced by education which helps them to develop greater confidence and the capacity to make decisions in regard to their own health.<sup>12</sup>

Noteworthy, is that marital status had no influence on birth preparedness. This however contradicts with a study conducted in north Ethiopia which showed women who were married were more likely to prepare for birth than the single women due to the cultural beliefs.<sup>13</sup>

## CONCLUSION

Birth preparedness is one of the components in Focused Antenatal Care that can be used to increase skilled deliveries in the country. As the International Community works towards Sustainable Development Goal of zero maternal deaths, there is need to target women in third trimester on birth preparedness to ensure risk of unskilled delivery is significantly reduced. There is need therefore for enhanced health education and awareness campaigns targeting women in their third trimester on birth preparedness.

## ACKNOWLEDGEMENTS

We would like to acknowledge the Department of Community Health and Development at Great Lakes University of Kisumu for the materials and continuous support in conducting this study. Our sincere gratitude goes to the County Director of Health and the Medical Superintendent Kirinyaga County for granting us the permission to carry out the study at Kirinyaga County. We also appreciate the support granted by the pregnant mothers and the staff working at the MCH/FP clinic during the study.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Great Lakes University Kisumu Research Ethics Committee (GREC)*

## REFERENCES

1. Hiluf M, Fantahum M. Birth Preparedness and Complication Readiness among women in north Ethiopia. *Ethiopian journal health development*. 2008;22:8-20.
2. World Health Organization. Birth and emergency preparedness in antenatal care: Integrated management of pregnancy and childbirth (IMPAC). Geneva, Switzerland, 2006.
3. World Health Organization. The World Health Report 2005: Make every mother and child count. Geneva, Switzerland, 2005.
4. Nahar S, Banu M, Nasreen HE. Women-focused development intervention reduces delays in accessing emergency obstetric care in urban slums in Bangladesh: a cross-sectional study. *BMC Pregnancy and Childbirth*. Available at: <http://www.biomedcentral.com/1471-2393/11/11>. Accessed on May, 2011.
5. Kenya National Bureau of Statistics (KNBS) and ICF Macro. 2010. Kenya Demographic and Health Survey 2008-09. Calverton, Maryland: KNBS and ICF Macro.
6. Kenya National Bureau of Statistics (KNBS) and ICF Macro. 2014. Kenya Demographic and Health Survey 2014/15. Calverton, Maryland: KNBS and ICF Macro.
7. Ministry of Health. National Road Map for Accelerating the Attainment of the MDGs Related to Maternal and Newborn Health in Kenya, 2010.
8. John Hopkins Program for International Education in Gynecology and Obstetrics (2010). Monitoring birth preparedness and complication readiness: Tools and indicators for maternal and newborn health. Baltimore, Maryland: Author.
9. Onah HE, Ikeako LC, Iloabachie GC. Factors associated with the use of maternity services in Enugu, southeastern Nigeria. *Social Sci Med J*. 2006;63:1870-8.
10. Hailu M, Gebremariam A, Alemseged F, Deribe K. Birth Preparedness and Complication Readiness among Pregnant Women in Southern Ethiopia. *PLoS One*. 2011;6(6):e21432.
11. Mpembeni RN, Killewo JZ, Leshabari MT, Massawe SN, Jahn A, Mushi D, et al. Use pattern of maternal health services and determinants of skilled care during delivery in Southern Tanzania: implications for achievement of MDG-5 targets. *BMC Pregnancy and Childbirth*. 2007;7(29):1-7.
12. Nigussie M, Mariam D, Mitike G. Assessment of safe delivery service utilization among women of child bearing age North Gondar Zone, North West Ethiopia. *Ethiopian J Health Development*. 2004;18:146-52.
13. Hiluf M, Fantahum M. Birth Preparedness and Complication Readiness among women in north Ethiopia. *Ethiopian J Health Development*. 2008;22:8-20.

**Cite this article as:** Karimi OW, Murigi MW, Pertet A, Odawa CO. Factors associated with level of birth preparedness among pregnant mothers at Kerugoya county referral hospital. *Int J Community Med Public Health* 2017;4:3998-4002.