Review Article

DOI: https://dx.doi.org/10.18203/2394-6040.ijcmph20212324

Guillain-Barré syndrome: pathophysiology, etiology, causes, and treatment

Rahmathulla S. Rahman^{1*}, Moayyad S. Bauthman², Amer M. Alanazi³, Naif N. Alsillah⁴, Ziyad M. Alanazi³, Mahdi I. Almuhaysin⁵, Ruyuf K. Almutairi⁶, Kadhem S. Binobaid⁵, Yousef H. Alharthi⁷, Rime M. Bawareth⁸, Omar N. Alrawili³

Received: 17 May 2021 Revised: 29 May 2021 Accepted: 31 May 2021

*Correspondence:

Dr. Rahmathulla S. Rahman, E-mail: safiyul@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Guillain—Barré syndrome (GBS) is a polyradiculoneuropathy autoimmune disease that is characterized by significant inflammation that affects the peripheral nervous system in a rapidly progressive pattern that is mainly clinically presented by muscle weakness. The present literature review aims to broadly discuss GBS: etiology, pathophysiology and management in order to gain an understading of the existing studies that are relevant to this literature review. Among the reported antibodies, anti-GM1 and anti-GQ1B have been reported to be responsible for attacking and damaging either the neuromuscular junctions or peripheral nerves. Moreover, it has been found that the anti-GD1a antibodies in patients bind to the neuromuscular junction and also bind to the nodes of Ranvier of the peripheral nerves and the paranodal myelin of the affected nerves. Reports have shown that this disease is identified as special forms of neuropathies that develop in immune-mediated, post-infection sequelae. Furthermore, in another study it was reported that Molecular mimicry has been previously reported to significantly correlate with the development of the disease as it was investigated in animal models. In addition, Campylobacter jejuni, a pathogen that causes gastrointestinal infections has been previously reported to predispose to the development of GBS in humans. However, scientists have found that plasma exchange and intravenous immunoglobulins (IVIG) remain the most significant and efficacious factors in managing the disease. Nevertheless, recent trials have investigated other approaches that are less efficacious and can lead to serious adverse events and complications.

Keywords: Guillain-Barré syndrome, Neurology, Management, Epidemiology, Etiology

INTRODUCTION

Guillain-Barré syndrome (GBS) is a polyradiculoneuropathy autoimmune disease that is characterized by significant inflammation that affects the peripheral nervous system in a rapidly progressive pattern that is mainly clinically presented by muscle weakness in addition to adjacent sensory pictures.¹ The symptoms can

¹Department of Internal Medicine, Sameera Medical Center, Jeddah, Saudi Arabia

²Department of Internal Medicine, ⁸College of Medicine, King Abdulaziz University Hospital, Jeddah, Saudi Arabia

³College of Medicine, Northern Border University, Arar, Saudi Arabia

⁴Department of Emergency Medicine, New Najran Hospital, Najran, Saudi Arabia

⁵College of Medicine, Medical University of Lodz, Lodz, Poland

⁶College of Medicine, Ibn Sina National College, Jeddah, Saudi Arabia

⁷College of Medicine, University of Tabuk, Tabuk, Saudi Arabia

⁸College of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia

develop acutely or sub acutely within few days to few weeks.² The severity of the disease is hugely variable and can be mild or very severe that may cause serious morbidities and even mortality, such as autonomic dysfunction, respiratory failure and tetraplegia. Estimates show that the death from GBS occurs 2-3% of the cases among patients within the developed countries.3,4 However, previous studies have demonstrated that intravenous immunoglobulins (IVIG) administration and performing plasma exchange can significantly enhance the prognosis of the disease and lead to recovery. 5,6 Although the prognosis of the disease tends to be good in many cases, it has been estimated that around 15% of the cases with GBS have poor outcomes, despite the adequate management with the currently-validated modalities owing to the development of severe complications especially respiratory muscle weakness, requiring patients to be mechanically-ventilated. Recent reports from Western countries have furtherly showed that the national campaigns aim to eliminate poliomyelitis as the GBS is now the leading cause of flaccid paralysis both the acute and subacute types.7 According to the various global reports, previous studies have revealed that the incidence of GBS rate between 0.16 and 0.4 /100,000 year in individual of all ages.^{2,8}

The diagnosis of GBS is mainly based on the clinical sequelae that patients usually have. This includes the development of lower limbs muscle weakness, reduced or absent deep reflexes and the monophasic pattern of the disease. Also, the nerve conductions studies and cerebrospinal fluid analysis are also important diagnostic tests that can help to confirm the diagnosis. 9,10 Moreover, the disease includes a number of variants that differ from each other in the pattern of the clinical presentations related to the severity and prognosis of the condition. Moreover, previous reports indicated that these variants also differ from each other in the laboratory findings and the pathophysiology of the disease. The present literature review aims to broadly discuss the GBS: etiology, pathophysiology and management in order to gain an understading of the existing studies that are relevant to this literature review.

METHODOLOGY

A systematic search was conducted to identify relevant studies in the following databases: PubMed, Medline, Web of Science, Embase, Google Scholar, and Scopus. The following search terms were used: "Guillain–Barré syndrome" or "Guillain–Barré", "pathophysiology" and "management" or "outcome" and "eitiology" or "etiology" or "aetiology". The reference lists were manually searched to identify additional relevant studies meeting inclusion criteria. We included any study that reports GBS and its pathophysiology, causes and outcomes. No restrictions were applied.

DISCUSSION

Etiology

Specific factors have been previously reported to correlate with the development of GBS and its variants. Reports have shown that the disease is identified as special forms of neuropathies that develop in immune-mediated, postinfection sequelae. Molecular mimicry has been previously reported to significantly correlate with the development of the disease as it was investigated in animal models. Campylobacter jejuni, a pathogen that causes gastrointestinal infections has been previously reported to predispose to the development of GBS in humans. This is probably due to the similar antigenicity between the gangliosides that are involved in the structure of the peripheral nerve cells and the lipooligosaccharide component of the outer membranous layer of the bacterium, leading to a similar antibody-mediated attack against these nerve cells.11 Accordingly, whenever the infection occurs, it is anticipated that the neuropathic sequelae of GBS will secondarily develop as an immunemediated reaction.

In addition to *Campylobacter jejuni*, other infections causing gastrointestinal and respiratory illnesses have also been correlated with the development of GBS. Within the first 1-6 weeks of eliciting GBS-related symptoms, it has been reported that more than two-thirds of the affected patients have been diagnosed with a previous infection. ¹⁰ For instance, Dirlikov et al previously reported that many cases of GBS have been reported and diagnosed by following the infection of the Zika virus during its relevant previous pandemic. ¹²

Moreover, previous cases have reported that some other etiologies other than the viral infections as some medications and surgical manipulations might also predispose to the development of GBS.¹³ In the early vaccination campaigns against influenza A/H1N1 by the flu vaccines in 1976, it was noticable that many cases of GBS and neuropathies have developed in relation to the vaccination administration, although it was reported that within the following years, the numbers of cases of GBS following the vaccination has significantly reduced to only one case per one million vaccines administered. In the same context, previous studies have demonstrated that GBS is seven times more likely to develop following influenza infection than being vaccinated against the disease.¹⁴⁻¹⁸

Epidemiology

The burden of GBS on the global healthcare facilities cannot be ignored. Despite that this disease is marked as a rare disorder that affects some minor populations, the estimated incidence for GBS varies from 0.4 and 0.2 per 100,000 population. A total of \$318,966 cost burden has been estimated for the provided medical care for patients with GBS. Moreover, previous reports also has shown that

the annual costs for taking care of GBS patients might reach a total of \$1.7 billion per year. Regarding gender, the estimation has shown that male patients are more frequently diagnosed with GBS than females. Besides, evidence from the current literature shows that GBS can affect up to 100,000 patients per year globally. 19-21

Pathophysiology

This section is extensively discussed studies related to this literature. Estimates show that infections contribute to the development of GBS as previous investigations have shown that in 70% of cases with GBS, patients had a history of infections.²² Accordingly, efforts were directed to understand more about the potential role and molecular similarity that the infecting organisms might have with the cells and tissues of the infected patients that developed GBS especially the axonal type. Yuki et al previously indicated this by showing that the molecular structures of the gangliosides that are found within the peripheral nerves are similar to the lipooligosaccharide that is extensively found in the structure of Campylobacter jejuni. 11 Previous studies have indicated this theory by laboratory experiments in rabbits that were subjected to the lipooligosaccharide that mimics the human gangliosides of peripheral nerves and found that these rabbits significantly developed GBS-like symptoms (acute motor axonal neuropathy) more specifically developing flaccid tetraplegia. 23,24

Antibodies that concomitantly develop against the host gangliosides attack different parts of the peripheral nervous system. Among the reported antibodies, GM1 and GQ1B antibodies have been reported to be responsible for attacking and damaging either neuromuscular junctions or peripheral nerves. ^{25,26} Moreover, it has been found that the anti-GD1a antibodies in patients bind to the neuromuscular junction and also bind to the nodes of Ranvier of the peripheral nerves and the paranodal myelin of the affected nerves. ^{27,28} According to the different parts that have been affected by the different antibodies and antigenic stimulation, it has been suggested that such differences are the major causes of the variation of the course of the disease.

Besides, a previous investigation by Susuki et al showed that after being infected with the pathogenic organism that presented with a similar antigenicity, complement activation was observed among the infected patients which was an indicator of its major role in the pathogenesis and development of the disease.²⁹ As previously mentioned, it has been found that certain antibodies might predispose to certain variants and clinical symptoms of GBS. For instance, anti-GQ1B antibodies have been previously reported to correlate with Miller-Fisher syndrome while it has been reported that anti-GM1 antibodies predispose to the development of the variant of axonal motor neuropathy.^{30,31} Moreover, anti-GT1A antibodies have also been reported to have a direct correlation with the development of the GBS variant pharyngeal-cervical-

brachial syndrome.³² However, the validity of these antibodies including both the specificity and sensitivity in detection and diagnosis of the different variants remains low and should be only considered for confirmation of the evaluation and diagnosis. Therefore, further investigations might be needed for further elucidation of the potential primary roles that these antibodies might play in the pathogenesis and differentiation of GBS and its variants, which will be reflected in inaugurating better diagnostic and management modalities of the disease. Additionally, although acute inflammatory demvelinating polyneuropathy has been marked as the most prevalent variant of GBS within the United States, the pathophysiology of this disorder remains vague and further investigations are needed.

Management

Previous randomized controlled trials have demonstrated the efficacy of previous modalities in the management of GBS. Plasma exchange and intravenous immunoglobulins (IVIG) remain the most efficacious and currently validated modalities that are recommended for administration in patients GBS. with Furthermore, the previous investigations have shown that plasma exchange is being useful by enhancing the removal of antibodies complement proteins and humoral mediators that are usually observed mainly contribute to the development and pathophysiology of GBS. It has been recommended that plasma exchange should be administered on five different sessions as a volume exchange. However, the exact mechanism of action of plasma exchange has not been fully understood yet and still needs further investigations. On the other hand, it has been observed that IVIG might have potent immune-modulating activities, although it has been reported that the main mechanism behind the action and physiology of IVIG in playing this role remains controversial, and has not been proven, yet. IVIG should be administered over five days and a total dose of 2 grams per kilogram should be considered.³³ However, previous investigations that compared the efficacy and safety of both modalities have demonstrated that both of them equally contribute to the successful management of the disease.34

Previous studies have demonstrated that for both modalities to be efficacious in the management of the disease. Any of them should be administered for consecutive four weeks with no advantage of any of them over the other. Besides, for a more efficacious and successful management plan, treatment with any of these modalities should be initiated within the first two weeks. 35-37 Although corticosteroids have been used in the management of immune-mediated reactions within the past decades, recent investigations have demonstrated that the efficacy of using corticosteroids (whether the intravenous or the oral combinations) was observed to be less than that of using placebo or when compared to the efficacy of using IVIG and plasma exchange whether alone or when used in combination. In general, it has been

reported that the applications of these management modalities can significantly decrease the recovery period from GBS. In previous investigations, independent ambulations were observed among patients that were treated with the previous management modalities within 32 days from the start of the disease course. 5,34,36,38-40

Reports have shown that the prognosis of GBS is generally good and estimates show that around 80% of patients and more have a significant chance of recovery and reduced morbidities; however, the other percentage of patients usually develop complications and significant morbidities. Moreover, the concurrent administration of plasma exchange and IVIG at time intervals or the administration of steroids following IVIG administration were not associated with significant favorable events. 41,42 This was indicated by a recent 2021 randomized controlled trial, which reported a second IVIG dose was not significantly associated with any significant improvements in patients with severe GBS and it can negatively impact the prognosis by inducing some adverse events.⁴³ The same results were also previously reported by an observational study by Verboon et al which showed that second IVIG doses could not significantly enhance the prognosis of poor GBS.⁴⁴ Although previous investigations have approached the efficacy of some other management modalities, none of these approaches have been proven to be significantly efficacious as compared to plasma exchange and IVIG.⁴⁵

CONCLUSION

In the present study, we have reviewed the findings from previous studies about the GBS: etiology, pathophysiology and management. Viral infections are the most frequent causes that predispose to the development of GBS. Therefore, proper interventions should be offered for the patients at risk. Plasma exchange and IVIG remain the most significant and efficacious factors in managing the disease. However, recent trials have investigated other approaches that have been found to be less efficacious and can lead to serious adverse events and complications.

Funding: No funding sources Conflict of interest: None declared Ethical approval: Not required

REFERENCES

- Hughes RA, Cornblath DR. Guillain-Barré syndrome. Lancet. 2005;366(9497):1653-66.
- 2. Cosi V, Versino M. Guillain-Barré syndrome. Neurol Sci. 2006;27(1):47-51.
- 3. Van den Berg B, Walgaard C, Drenthen J, Fokke C, Jacobs BC, van Doorn PA. Guillain-Barré syndrome: pathogenesis, diagnosis, treatment and prognosis. Nat Rev Neurol. 2014;10(8):469-82.
- 4. Dornonville de la Cour C, Jakobsen J. Residual neuropathy in long-term population-based follow-up of Guillain-Barré syndrome. Neurology. 2005;64(2):246-53.

- Hughes RA, Swan AV, van Doorn PA. Intravenous immunoglobulin for Guillain-Barré syndrome. Cochrane Database Systemat Rev. 2014;9:CD002063.
- 6. Raphaël JC, Chevret S, Hughes RA, Annane D. Plasma exchange for Guillain-Barré syndrome. Cochrane Database Systemat Rev. 2002(2):CD001798.
- 7. Olivé JM, Castillo C, Castro RG, de Quadros CA. Epidemiologic study of Guillain-Barré syndrome in children <15 years of age in Latin America. J Infect Dis. 1997;175(1):160-4.
- 8. Govoni V, Granieri E. Epidemiology of the Guillain-Barré syndrome. Curr Opinion Neurol. 2001;14(5):605-13.
- 9. Asbury AK, Cornblath DR. Assessment of current diagnostic criteria for Guillain-Barré syndrome. Ann Neurol. 1990;27:21-4.
- Fokke C, van den Berg B, Drenthen J, Walgaard C, van Doorn PA, Jacobs BC. Diagnosis of Guillain-Barré syndrome and validation of Brighton criteria. Brain J Neurol. 2014;137(1):33-43.
- 11. Yuki N, Taki T, Inagaki F. A bacterium lipopolysaccharide that elicits Guillain-Barré syndrome has a GM1 ganglioside-like structure. J Exp Med. 1993;178(5):1771-5.
- 12. Dirlikov E, Major CG, Medina NA. Clinical Features of Guillain-Barré Syndrome With vs Without Zika Virus Infection, Puerto Rico, 2016. JAMA Neurol. 2018;75(9):1089-97.
- 13. Awong IE, Dandurand KR, Keeys CA, Maung-Gyi FA. Drug-associated Guillain-Barré syndrome: a literature review. Ann Pharmacother. 1996;30(2):173-80.
- 14. Yuki N, Hartung HP. Guillain-Barré syndrome. N Engl J Med. 2012;366(24):2294-304.
- 15. Campbell AM. The aetiology of polyneuritis. Proc Royal Soc Med. 1958;51(3):157-9.
- 16. Cao-Lormeau VM, Blake A, Mons S. Guillain-Barré Syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. Lancet. 2016;387(10027):1531-9.
- 17. Schonberger LB, Bregman DJ, Sullivan-Bolyai JZ. Guillain-Barre syndrome following vaccination in the National Influenza Immunization Program, United States, 1976--1977. Am J Epidemiol. 1979;110(2):105-23.
- Lasky T, Terracciano GJ, Magder L. The Guillain-Barré syndrome and the 1992-1993 and 1993-1994 influenza vaccines. N Engl J Med. 1998;339(25):1797-802.
- 19. Willison HJ, Jacobs BC, van Doorn PA. Guillain-Barré syndrome. Lancet. 2016;388(10045):717-27.
- 20. Sejvar JJ, Baughman AL, Wise M, Morgan OW. Population incidence of Guillain-Barré syndrome: a systematic review and meta-analysis. Neuroepidemiology. 2011;36(2):123-33.
- 21. Nguyen TP, Taylor RS. Guillain Barre Syndrome. In: StatPearls. Treasure Island (FL): StatPearls Publishing. StatPearls Publishing LLC. 2021.

- 22. Jacobs BC, Rothbarth PH, van der Meché FG. The spectrum of antecedent infections in Guillain-Barré syndrome: a case-control study. Neurology. 1998;51(4):1110-5.
- 23. Susuki K, Nishimoto Y, Yamada M. Acute motor axonal neuropathy rabbit model: immune attack on nerve root axons. Ann Neurol. 2003;54(3):383-8.
- 24. Yuki N, Yamada M, Koga M. Animal model of axonal Guillain-Barré syndrome induced by sensitization with GM1 ganglioside. Ann Neurol. 2001;49(6):712-20.
- 25. Willison HJ, O'Hanlon G, Paterson G, et al. Mechanisms of action of anti-GM1 and anti-GQ1b ganglioside antibodies in Guillain-Barré syndrome. J Infect Dis. 1997;176(2):144-9.
- 26. Greenshields KN, Halstead SK, Zitman FM. The neuropathic potential of anti-GM1 autoantibodies is regulated by the local glycolipid environment in mice. J Clin Investig. 2009;119(3):595-610.
- Chiba A, Kusunoki S, Obata H, Machinami R, Kanazawa I. Serum anti-GQ1b IgG antibody is associated with ophthalmoplegia in Miller Fisher syndrome and Guillain-Barré syndrome: clinical and immunohistochemical studies. Neurology. 1993;43(10):1911-7.
- 28. Goodfellow JA, Bowes T, Sheikh K. Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy. J Neurosci. 2005;25(7):1620-8.
- 29. Susuki K, Rasband MN, Tohyama K. Anti-GM1 antibodies cause complement-mediated disruption of sodium channel clusters in peripheral motor nerve fibers. J Neurosci. 2007;27(15):3956-67.
- 30. Chiba A, Kusunoki S, Shimizu T, Kanazawa I. Serum IgG antibody to ganglioside GQ1b is a possible marker of Miller Fisher syndrome. Ann Neurol. 1992;31(6):677-9.
- 31. Gregson NA, Jones D, Thomas PK, Willison HJ. Acute motor neuropathy with antibodies to GM1 ganglioside. J Neurol. 1991;238(8):447-51.
- 32. O'Leary CP, Veitch J, Durward WF, Thomas AM, Rees JH, Willison HJ. Acute oropharyngeal palsy is associated with antibodies to GQ1b and GT1a gangliosides. J Neurol Neurosurg Psychiatry. 1996;61(6):649-51.
- 33. Ortiz-Salas P, Velez-Van-Meerbeke A, Galvis-Gomez CA, Rodriguez QJ. Human Immunoglobulin Versus Plasmapheresis in Guillain-Barre Syndrome and Myasthenia Gravis: A Meta-Analysis. J Clin Neuromuscular Dis. 2016;18(1):1-11.
- Van der Meché FG, Schmitz PI. A randomized trial comparing intravenous immune globulin and plasma exchange in Guillain-Barré syndrome. Dutch Guillain-Barré Study Group. N Engl J Med. 1992;326(17):1123-9.
- 35. Raphaël JC, Chevret S, Hughes RA, Annane D. Plasma exchange for Guillain-Barré syndrome. The

- Cochrane Database Systemat Rev. 2012(7):CD001798.
- 36. Plasmapheresis and acute Guillain-Barré syndrome. The Guillain-Barré syndrome Study Group. Neurology. 1985;35(8):1096-104.
- 37. Appropriate number of plasma exchanges in Guillain-Barré syndrome. The French Cooperative Group on Plasma Exchange in Guillain-Barré Syndrome. Ann Neurol. 1997;41(3):298-306.
- 38. Hughes RA. Plasma exchange versus intravenous immunoglobulin for Guillain-Barré syndrome. Therapeutic Apheresis. 1997;1(2):129-30.
- 39. Hughes RA, Brassington R, Gunn AA, van Doorn PA. Corticosteroids for Guillain-Barré syndrome. Cochrane Database Systemat Revi. 2016;10(10):CD001446.
- 40. McKhann GM, Griffin JW, Cornblath DR, Mellits ED, Fisher RS, Quaskey SA. Plasmapheresis and Guillain-Barré syndrome: analysis of prognostic factors and the effect of plasmapheresis. Ann Neurol. 1988;23(4):347-53.
- 41. Van Koningsveld R, Schmitz PI, Meché FG, Visser LH, Meulstee J, van Doorn PA. Effect of methylprednisolone when added to standard treatment with intravenous immunoglobulin for Guillain-Barré syndrome: randomised trial. Lancet. 2004;363(9404):192-6.
- 42. Randomised trial of plasma exchange, intravenous immunoglobulin, and combined treatments in Guillain-Barré syndrome. Plasma Exchange/Sandoglobulin Guillain-Barré Syndrome Trial Group. Lancet. 1997;349(9047):225-30.
- 43. Walgaard C, Jacobs BC, Lingsma HF, et al. Second intravenous immunoglobulin dose in patients with Guillain-Barré syndrome with poor prognosis (SID-GBS): a double-blind, randomised, placebocontrolled trial. Lancet Neurol. 2021;20(4):275-83.
- 44. Verboon C, van den Berg B, Cornblath DR. Original research: Second IVIg course in Guillain-Barré syndrome with poor prognosis: the non-randomised ISID study. J Neurol Neurosurg Psychiatr. 2020;91(2):113-21.
- 45. Doets AY, Hughes RA, Brassington R, Hadden RD, Pritchard J. Pharmacological treatment other than corticosteroids, intravenous immunoglobulin and plasma exchange for Guillain-Barré syndrome. Cochrane Database Systemat Revi. 2020;1(1):CD008630.

Cite this article as: Rahman RS, Bauthman MS, Alanazi AM, Alsillah NN, Alanazi ZM, Almuhaysin MI, et al. Guillain—Barré syndrome: pathophysiology, etiology, causes, and treatment. Int J Community Med Public Health 2021;8:3624-8.